

COLONOSCOPY PREP WITH MOVIPREP

CLEAR LIQUID DIET

THE DAY BEFORE YOUR PROCEDURE YOU WILL BE ON A CLEAR LIQUID DIET ALL DAY!! THIS INCLUDES APPLE JUICE, GATORADE, CHICKEN BROTH, JELL-0, POPSICLES, AND VANILLA ICE CREAM, NOTHING RED. BLUE OR PURPLE. COFFEE AND TEA ARE CONSIDERED A CLEAR LIQUID SO THEY ARE OK.


***THE INSTRUCTIONS ARE ATTACHED ON THE SECOND PAGE. PLEASE CALL US IF YOU HAVE ANY QUESTIONS REGARDING YOUR PREP. MAKE SURE YOU HAVE PLENTY OF CLEAR LIQUIDS THROUGHOUT THE DAY. AND YOU CAN NOT HAVE ANYTHING MORE AFTER 12 MIDNIGHT. AGAIN, CALL THE OFFICE IF YOU HAVE ANY QUESTIONS (904) 388-8686.

PATIENT INSTRUCTIONS EVENING-ONLY DOSING

The MoviPrep® carton contains 4 pouches and a disposable container for mixing.
You must complete the entire prep to ensure the most effective cleansing.

BEGIN FIRST DOSE AT _____


BEGIN SECOND DOSE AT _____



STEP 1: MIX DOSE

- › Empty 1 Pouch A and 1 Pouch B into the disposable container
- › Add lukewarm drinking water to the top line of the container and mix

NOTE: You can mix solution ahead of time and refrigerate prior to drinking.
The reconstituted solution should be used within 24 hours.



STEP 2: DRINK DOSE

- › The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is consumed



STEP 3: DRINK CLEAR LIQUIDS

- › **Drink 16 oz of the clear liquid of your choice.** This is a necessary step to ensure adequate hydration and an effective prep

Clear liquids include: water, clear fruit juices without pulp, including apple, white grape, or white cranberry, strained limeade or lemonade, coffee or tea (do not use dairy or non-dairy creamer), clear broth, clear soda, gelatin (without added fruit or topping), and popsicles (without pieces of fruit or fruit pulp). Ask your doctor if you have any questions about whether a particular drink is acceptable.

Follow steps 1-3 to complete first dose. Repeat steps 1-3 for second dose.

If you have any questions, please call our office at _____

A colonoscopy prep causes the body to lose a significant amount of fluid and can result in sickness due to dehydration.
Stay hydrated by drinking all required clear liquids during the prep.

IMPORTANT SAFETY INFORMATION (CONT'D)

If you have a condition that destroys red blood cells called glucose-6-phosphate dehydrogenase (G6PD) deficiency, talk to your doctor before taking MOVIPREP.

Please be advised that MOVIPREP contains 131 mg of phenylalanine per treatment.

If you are pregnant or planning to become pregnant, talk to your doctor before taking MOVIPREP.

In clinical studies with MOVIPREP, the most common side effects for split dosing were malaise (feeling uneasy), nausea, stomach (abdominal) pain, vomiting, and upper stomach pain. The most common side effects for evening only dosing were stomach swelling (abdominal distension), anal discomfort, thirst, nausea, stomach (abdominal) pain, sleep disorder, rigors (body shakes), hunger, malaise (feeling uneasy), vomiting, and dizziness.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch/ or call 1-800-FDA-1088.

For product information, adverse event reports, and product complaint reports, please contact:

Salix Product Information Call Center:

Phone: 1-800-508-0024 | Fax: 1-510-595-8183 | Email: salix@medcomsol.com

Please see complete Prescribing Information at MOVIPREP.com.



Web site: MoviPrep.com 8510 Colonnade Center Drive, Raleigh, NC 27615 Tel: 866.669.SLXP (7597).
MoviPrep® is a registered trademark and PM/JAM Split Dosing™ is a trademark of Salix Pharmaceuticals, Inc.
© 2014 Salix Pharmaceuticals, Inc. All rights reserved. Printed in USA. MOV5-0114

MoviPrep

(PEG-3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate, and ascorbic acid for oral solution, 100 g/7.5 g/2.691 g/1.015 g/5.9 g/4.7 g)